




It's the way we're *wired*

GRADE 11 MATHS

Charmaine



FUNCTIONS

Functions of the general form $y = ax^2 + bx + c$ are called parabolic functions. The constants a , b and c have different effects on the parabola

The effect of a


The sign of a determines the shape of the graph.

- For $a > 0$, the graph of $f(x)$ is a “smile”. The graph of $f(x)$ is stretched vertically upwards; as a gets larger, the graph gets narrower.

For $0 < a < 1$, as a gets closer to 0, the graph of $f(x)$ gets wider.

- For $a < 0$, the graph of $f(x)$ is a “frown”. The graph of $f(x)$ is stretched vertically downwards; as a gets smaller, the graph gets narrower.

For $-1 < a < 0$, as a gets closer to 0, the graph of $f(x)$ gets wider




FUNCTIONS

The effect of b:

The x-value of the turning point is determined by $x = -\frac{b}{2a}$. This is the axis of symmetry

The effect of c:

c gives us the y intercept i.e. (0;c)



EXAMPLE: FUNCTIONS


Sketch the graph of $y = \frac{1}{2}x^2 - 4x + \frac{7}{2}$

Determine the intercepts, turning point and the axis of symmetry. Give the domain and range of the function.

We notice that $a = \frac{1}{2}$; $b = -4$ $c = \frac{7}{2}$

Step 1: Examine the equation of the form $y = ax^2 + bx + c$

We notice that $a > 0$, therefore the graph is a “smile” and has a minimum turning point.



EXAMPLE: FUNCTIONS

Step 2: Determine the turning point and the axis of symmetry

$$x = -\frac{b}{2a}$$


$$x = -\frac{-4}{2\left(\frac{1}{2}\right)} = 4$$

Therefore the axis of symmetry is $x = 4$.

Substitute $x = 4$ into the original equation to obtain the corresponding y -value.

$$y = \frac{1}{2}(4)^2 - 4(4) + \frac{7}{2} = -4\frac{1}{2}$$

This gives the point $\left(4; -4\frac{1}{2}\right)$ as the turning point



EXAMPLE: FUNCTIONS

Step 3: Determine the y-intercept

The y-intercept is obtained by letting $x = 0$: $y = \frac{7}{2}$

This gives the point $(0; \frac{7}{2})$ as the y-intercept

Step 4: Determine the x-intercepts

The x-intercepts are obtained by letting $y = 0$:

$$\begin{aligned} 0 &= \frac{1}{2}x^2 - 4x + \frac{7}{2} \\ &= x^2 - 8x + 7 \\ &= (x - 7)(x - 1) \end{aligned}$$


Either $x = 7$ or 1

This gives the points $(1; 0)$ and $(7; 0)$.



EXAMPLE: FUNCTIONS


- Step 5: Plot the points and sketch the graph



EXAMPLE: FUNCTIONS

Step 6: State the domain and range

- Domain: $\{x: x \in R\}$
- Range: $\{y: y \geq -4\frac{1}{2}, y \in R\}$



EXERCISE: FUNCTIONS

Sketch graphs of the following functions and determine:


- intercepts
- turning point
- axes of symmetry
- domain and range

a) $y = -x^2 + 4x + 5$

b) $y = 2(x + 1)^2$


c) $y = 3x^2 - 2(x + 2)$

d) $y = 3(x - 2)^2 + 1$



SOLUTIONS: FUNCTIONS

a)



Intercepts: $(-1; 0)$, $(5; 0)$, $(0; 5)$

Turning point: $(2; 9)$


Axes of symmetry: $x = 2$

Domain: $\{x : x \in \mathbb{R}\}$

Range: $\{y : y \leq 9, y \in \mathbb{R}\}$

SOLUTIONS: FUNCTIONS

b)



Intercepts: $(-1; 0), (0; 2)$

Turning point: $(-1; 0)$


Axes of symmetry: $x = -1$

Domain: $\{x : x \in \mathbb{R}\}$

Range: $\{y : y \geq 0, y \in \mathbb{R}\}$

SOLUTIONS: FUNCTIONS

c)



Intercepts: $(-0,87; 0)$, $(1,54; 0)$, $(0; -4)$

Turning point: $(0,33; -4,33)$

Axes of symmetry: $x = -0,33$


Domain: $\{x : x \in \mathbb{R}\}$

Range: $\{y : y \geq 4,33, y \in \mathbb{R}\}$



SOLUTIONS: FUNCTIONS

d)



Intercepts: $(0; 13)$ Turning point: $(2; 1)$

Axes of symmetry: $x = 2$

Domain: $\{x : x \in \mathbb{R}\}$


Range: $\{y : y \geq 1, y \in \mathbb{R}\}$



FUNCTIONS: FINDING THE EQUATION OF A PARABOLA

If the intercepts are given, use $y = a(x - x_1)(x - x_2)$.

Example:



x -intercepts: $(-1; 0)$ and $(4; 0)$

$$y = a(x - x_1)(x - x_2)$$

$$= a(x + 1)(x - 4)$$

$$= ax^2 - 3ax - 4a$$

y -intercept: $(0; 2)$

$$-4a = 2$$

$$a = -\frac{1}{2}$$

Equation of the parabola:

$$y = ax^2 - 3ax - 4a$$


$$= -\frac{1}{2}x^2 - 3\left(-\frac{1}{2}\right)x - 4\left(-\frac{1}{2}\right)$$

$$= -\frac{1}{2}x^2 + \frac{3}{2}x + 2$$

FUNCTIONS: FINDING THE EQUATION OF A PARABOLA

If the x -intercepts and another point are given, use $y = a(x - x_1)(x - x_2)$.

Example:



x -intercepts: $(1; 0)$ and $(5; 0)$

$$y = a(x - x_1)(x - x_2)$$

$$= a(x - 1)(x - 5)$$

$$= ax^2 - 6ax + 5a$$

Substitute the point: $(-1; 12)$

$$12 = a(-1)^2 - 6a(-1) + 5a$$

$$12 = a + 6a + 5a$$

$$12 = 12a$$

$$1 = a$$

Equation of the parabola:

$$y = ax^2 - 6ax + 5a$$


$$= x^2 - 6x + 5$$



FUNCTIONS: FINDING THE EQUATION OF A PARABOLA

If the turning point and another point are given, use $y = a(x + p)^2 + q$.

Example:



Turning point: $(-3; 1)$

$$y = a(x + p)^2 + q$$

$$= a(x + 3)^2 + 1$$

$$= ax^2 + 6ax + 9a + 1$$

Substitute the point: $(1; 5)$

$$5 = a(1)^2 + 6a(1) + 9a + 1$$

$$4 = 16a$$

$$\frac{1}{4} = a$$

Equation of the parabola:


$$y = \frac{1}{4}(x + 3)^2 + 1$$



EXERCISE: FUNCTIONS


Determine the equations of the following graphs

1.




EXERCISE: FUNCTIONS

2.




EXERCISE: FUNCTIONS

3.



EXERCISE: FUNCTIONS




SOLUTIONS: FUNCTIONS

1. $y = -3(x + 1)^2 + 6$ or $y = -3x^2 - 6x + 3$

2. $y = \frac{1}{2}x^2 - \frac{5}{2}x$

3. $y = \frac{2}{3}(x + 2)^2$

4. $y = -x^2 + 3x + 4$



FUNCTIONS: INVERSE

Given $f(x) = 2x^2$. Determine the inverse of $f(x)$. Sketch $f^{-1}(x)$ and $y = x$ on the same axes as $f(x)$


Step 1: Swap the x and y

$$y = 2x^2 \text{ becomes } x = 2y^2$$

Step 2: Make y the subject of the formula



$$y^2 = \frac{1}{2}x$$

$$y = \pm \sqrt{\frac{1}{2}x}$$



FUNCTIONS: INVERSE

- Not all inverses of functions are also functions. Some inverses of functions are relations. If an inverse is not a function, then we can restrict the **domain** of the **function** in order for the inverse to be a function.



FUNCTIONS: INVERSE

- To check for a function, draw a vertical line. If any vertical line cuts the graph in only one place, the graph is a function. If any vertical line cuts the graph in more than one place, then the graph is not a function.
- To check for a one-to-one function, draw a horizontal line. If any horizontal line cuts the graph in only one place, the graph is a one-to-one function. If any horizontal line cuts the graph in more than one place, then the graph is a many-to-one function.

